# Correct image plate positioning with the VistaPosition PSP Holder System





To preserve the image plate and for optimum image quality, we recommend using an original Dürr Dental light protection cover.

#### • Bitewing exposures

- AAAAAAAAAAAA 2º
- Position the image plate between the tongue and dental arch. Align the imaging plate orthoradially to avoid interdental space overlap.
- Actively guide the bite block downwards until it rests on the row of teeth.
- Close the mouth
- Position the aiming ring as close to the cheek as possible.
- Align the X-ray tube parallel to the indicator arm.
- Align the collimator according to the position of the image plate.




2° 🕅

ူမ္မ 🐺

### **3.** Lower jaw posterior teetth exposures



- Position the image plate between the tongue and dental arch. Actively guide the bite block downwards until it rests on the row of teeth and ensure that the patient holds it firmly after closing
- the mouth. • Let the indicator arm protrude horizontally from the mouth.
- Position the aiming ring as close to the cheek as possible.
- Align the X-ray tube parallel to the indicator arm.
- Align the collimator according to the position of the image plate.



## **5.** Lower jaw anterior teeth exposures



- Place the lower edge of the image plate at the base of the tongue behind the anterior region.
- Tilt the bite block to the cutting edge of the tooth to be captured.
- Have the patient push the lower jaw forward when closing. Actively guide the indicator arm upwards so that it protrudes
- from the mouth as horizontally as possible. • Position the aiming ring as close to the cheek as possible.
- Align the X-ray tube parallel to the indicator arm.
- Align the collimator according to the imaging plate position.



1<sup>1</sup>Ω 88 back

HANDON HAND HAND

#### 7. Lower jaw canine exposures



• Position the lower edge of the image plate at the base of the tongue behind the anterior teeth.

#### More tips & tricks and the electronic operating instructions can be found at www.duerrdental.com/intraoral-positioning

APPARARAA AAAAAAA



2°

**1**<sup>IQ</sup> 550 back

#### **2.** Endo exposures



# **O.** Upper jaw anterior teeth exposures



- Place the upper edge of the image plate on a sufficiently wide
- Tilt the bite block to the cutting edge of the tooth to be captured.
- Have the patient push the lower jaw forward when closing.
- Position the aiming ring as close to the cheek as possible.
- Align the X-ray tube parallel to the indicator arm.
- Align the collimator according to the position of the image plate.



#### 8. Upper jaw canine exposures



 Place the upper edge of the image plate on a sufficiently wide part of the palate.





- Turn the holder until the bite block is over the tip of the canine.
- Have the patient push the lower jaw towards the canine when closing.
- During closing, actively guide the indicator arm upwards so that it protrudes from the mouth as horizontally as possible.
- Position the aiming ring as close to the cheek as possible.
- Align the X-ray tube parallel to the indicator arm.
- Align the collimator according to the imaging plate position.





- Have the patient push the lower jaw towards the canine when closing.
- Position the aiming ring as close to the cheek as possible.
- Align the X-ray tube parallel to the indicator arm.
- Align the collimator according to the position of the image plate.



DÜRR DENTAL SE Höpfigheimer Str. 17 74321 Bietigheim-Bissingen Germany www.duerrdental.com info@duerrdental.com

P007100228L02/S04 Subject to technical alterations

